

SCIENTIFIC PAPERS
SERIES A. AGRONOMY
VOLUME LXI, No. 1, 2018

UNIVERSITY OF AGRONOMIC SCIENCES
AND VETERINARY MEDICINE OF BUCHAREST
FACULTY OF AGRICULTURE

SCIENTIFIC PAPERS
SERIES A. AGRONOMY

VOLUME LXI, No. 1

2018
BUCHAREST

SCIENTIFIC COMMITTEE

- Sinisa BERJAN – University of East Sarajevo, Bosnia and Herzegovina
- Dimitrios BILALIS – Agricultural University of Athens, Greece
- Iovu-Adrian BIRIȘ – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Lancelot BUTTERS – University of Central Lancashire, United Kingdom
- Raimundo CABRERA – University of La Laguna, Phytopathology Unit, Spain
- Costică CIONTU – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Sorin Mihai CÎMPEANU – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Stelica CRISTEA – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Ionela DOBRIN – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Marin DUMBRAVĂ – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Mihail DUMITRU – Research and Development Institute for Soil Science, Agro-chemistry and Environmental Protection of Bucharest, Romania
- Lenuța Iuliana EPURE – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Senol Zafer ERDOGAN – Konya Food and Agriculture University, Turkey
- André FALISSE – University of Liège, Gembloux Agro-Bio Tech, Belgium
- Cristian HERA – Romanian Academy
- Beatrice-Michaela IACOMI – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Cristian IACOMI – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Leonard ILIE - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Florin IMBREA – “King Mihai I of Romania” Banat University of Agricultural Sciences and Veterinary Medicine of Timișoara, Romania
- Viorel ION – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Mohsen JANMOHAMMADI – University of Maragheh, East Azarbaijan, Iran
- Gheorghe JIGĂU – State University of Moldova, Republic of Moldova
- Gerard JITĂREANU – “Ion Ionescu de la Brad” University of Agricultural Sciences and Veterinary Medicine of Iași, Romania
- Maria JOIȚA-PĂCUREANU – National Agricultural Research and Development Institute Fundulea, Romania
- Yalcin KAYA – Trakya University, Plant Breeding Research Center, Turkey
- Doru-Ioan MARIN – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Mircea MIHALACHE – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Mihai NICOLESCU – „Gheorghe Ionescu-Șișești” Academy of Agricultural and Forestry Sciences, Romania
- Ioan PĂCURAR – University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Romania
- Aurelian PENESCU – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Françoise PICARD-BONNAUD – University of Angers, France
- Teodor ROBU – „Ion Ionescu de la Brad” University of Agricultural Sciences and Veterinary Medicine of Iași, Romania
- Gheorghe Valentin ROMAN – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Mihail RURAC – State Agrarian University of Moldova, Republic of Moldova
- Teodor RUSU – University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Romania
- Dumitru Ilie SÂNDOIU – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Philippe SIMONEAU – University of Angers, France
- Gheorghe SIN – „Gheorghe Ionescu-Șișești” Academy of Agricultural and Forestry Sciences, Romania
- Vasilica STAN – University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Marin ȘTEFAN - University of Craiova, Romania
- David C. WEINDORF – Texas Tech University, USA
- Hristina YANCHEVA – Agricultural University of Plovdiv, Bulgaria

EDITORIAL BOARD

General Editor: Costică CIONTU

Executive Editor: Lenuța Iuliana EPURE

Members: Adrian Gheorghe BĂȘA, André FALISSE, Leonard ILIE, Viorel ION, Gheorghe JIGĂU, Doru Ioan MARIN, Mircea MIHALACHE

PUBLISHER: University of Agronomic Sciences and Veterinary Medicine of Bucharest,
Faculty of Agriculture, Romania

Address: 59 Mărăști Blvd, District 1, 011464, Bucharest, Romania

Phone/Fax: + 40 213 318-0466; E-mail: journal@agro-bucuresti.ro

Webpage: <http://agronomyjournal.usamv.ro>

Copyright 2018

To be cited: Scientific Papers. Series A. Agronomy, Vol. LXI, No. 1, 2018

The publisher is not responsible for the opinions published in the Volume. They represent the authors' point of view.

ISSN 2285-5785; ISSN CD-ROM 2285-5793; ISSN Online 2285-5807; ISSN-L 2285-5785

International Database Indexing: CABI, Index Copernicus, Google Scholar, CNCSIS B+, Ulrich's Periodicals Directory, Research Bible, Scipio, Scientific Indexing Service, PBN (Polish Scholarly Bibliography), OCLC (WorldCat); Web of Science Core Collection (Emerging Sources Citation Index)

SUMMARY

SOIL SCIENCES

1. TILLAGE EFFECTS ON SOIL ORGANIC CARBON, MICROBIAL BIOMASS CARBON AND BETA-GLUCOSIDASE ENZYME ACTIVITY IN A TYPIC HAPLOXERERT SOIL - **Mert ACAR, İsmail ÇELİK, Hikmet GÜNAL, Nurullah ACİR, Zeliha BARUT BEREKET, Mesut BUDAK** 13
2. THE EFFECTS OF DIFFERENT ZN DOSES AND MYCORRHIZAE APPLICATION ON HORSE BEAN GROWTH AND NUTRIENT UPTAKE UNDER STERILE AND NON STERILE SOIL CONDITIONS - **Cagdas AKPINAR, Ibrahim ORTAS, Ahmet DEMIRBAS** 21
3. CHISEL PLOW TILLAGE DEPTH EFFECT ON SOIL CARBON DIOXIDE EMISSION - **Ghassan AL-AZZAWI, Davut AKBOLAT** 27
4. EFFECT OF DIFFERENT MAGNESIUM DOSES ON GROWTH AND YIELD OF PEPPER PLANT IN MYCORRHIZA INOCULATED HARRAN SOIL - **Ahmet ALMACA** 34
5. SOIL EROSION RISK MAPPING USING USLE/GIS METHODOLOGY IN ROZE-CHAY CATCHMENT, NORTHWEST IRAN - **Farrokh ASADZADEH, Maryam RAHMATI, Hossein ASGARZADEH** 38
6. EVOLUTION OF DARK CHESTNUT STEPPE SOIL UNDER CONDITIONS OF DIFFERENT USE AND CLIMATE CHANGE - **Sveatoslav BALIUK, Lyudmila VOROTYNTSEVA** 42
7. INFLUENCE OF SOIL TILLAGE SYSTEMS AND INOCULATION ON SOYBEAN NODULATION AND YIELD - **Nicoleta CĂPĂȚÂNĂ, Ciprian BOLOHAN, Cristina Andreea OPREA, Doru Ioan MARIN** 46
8. AGROTECHNICAL SYSTEMS TO CONSERVING WATER IN THE SOIL FOR WHEAT CROP - **Felicia CHETAN, Teodor RUSU, Cornel CHETAN, Paula Ioana MORARU, Alina SIMON** 53
9. HEAVY METALS CONTENT IN ALFALFA CULTIVATED ON VERTISOLS ALONG THE HIGHWAY E75 FROM BELGRADE TO LESKOVAC (SERBIA) - **Zoran DINIĆ, Radmila PIVIĆ, Jelena MAKSIMOVIĆ, Aleksandar STANOJKOVIĆ, Dragana JOŠIĆ, Aleksandra STANOJKOVIĆ-SEBIĆ** 63
10. THE EFFECTS OF DIFFERENT SALT DOSES ON YIELD AND NUTRIENT UPTAKE OF TOMATO PLANT - **Hasan DURUKAN, Ahmet DEMIRBAS** 71
11. COMPARING THE EFFECTS OF COMPOST AND VERMICOMPOST ON CORN GROWTH, NUTRIENT CONCENTRATION AND UPTAKE DURING THE DIFFERENT GROWTH PERIODS - **İbrahim ERDAL, Ahmet DOGAN, Cennet YAYLACI, Pelin ALABOZ** 77
12. EVOLUTION MICROMORPHOLOGY OF ARGILLIC HORIZONS IN SOME ARID SOILS IN THE WEST OF URMIA LAKE IN WESTERN AZERBAIJAN PROVINCE, IRAN - **Tina JAHAN, Shahram MANAFI** 84

13. PHYSICAL PROPERTIES FEATURES OF ALLUVIAL IRRIGATED SOILS OF DNIESTER AND DNEPER RIVERS BASINS - Tamara LEAH, Valerian CERBARI, Sveatoslav BALIUK, Marina ZAKHAROVA, Oleksandr NOSONENKO	90
14. IMPACT OF CLIMATE CHANGE ON AGRO-CLIMATIC INDICATORS IN TRANSYLVANIAN PLAIN BETWEEN 2009-2016 - Mihai-Avram MAXIM, Teodor RUSU, Paula Ioana MORARU, Marius SĂBĂDAȘ, Ovidiu MAXIM	96
15. HEAVY METALS FROM THE SOIL AND MINERAL FERTILIZATION - Nicoleta MĂRIN, Nicoleta VRÎNCEANU, Naliana LUPAȘCU, Mihail DUMITRU	101
16. DYNAMICS OF SOIL PROPERTIES UNDER A POLLUTION GRADIENT IN URBAN AREAS (PLOVDIV, BULGARIA) - Slaveya PETROVA, Bogdan NIKOLOV, Iliana VELCHEVA, Mariya Yankova, Emiliya Kogan, Elena ZHELEVA, Ekaterina VALCHEVA, Alexandar ALEXANDROV, Mariana MARHOVA, Marinela TSANKOVA, Ivan ILIEV	109
17. THE USE OF CHEMICAL AND ORGANIC FERTILIZERS FOR SUNFLOWER CULTURE ON A STERILE DUMP - CĂTĂLIN AURELIAN ROȘCULETE, ELENA ROȘCULETE ...	117
18. INFLUENCE OF SYSTEM TILLAGE UPON WEEDING LERVELIN THE MAIZE CROP GROWN AT MOARA DOMNEASCĂ – ILFOV - Ionuț Cosmin SFETCU, Elena Loredana SFETCU, Doru Ioan MARIN	125
19. MYCORRHIZAS AS A TOOL IN MAPPING AGRICULTURAL SOILS - Vlad STOIAN, Roxana VIDICAN, Ioan ROTAR, Florin PĂCURAR, Mignon ȘANDOR, Mihai BUTA, Valentina STOIAN	130

CROP SCIENCES

1. MANAGEMENT OF FUSARIUM WILT (<i>Fusarium oxysporum</i> F. sp. <i>lycopersici</i>) OF TOMATO WITH ORGANIC AMENDMENTS - Gurama ABUBAKAR UMAR, Haruna SALISU GOMBE	139
2. EFFECT OF RHIZOBIUM BACTERIA ON NITROGEN FERTILIZER REQUEST OF COMMON VETCH (<i>Vicia sativa</i> L.) - Ayşen AKAY, Ali ÇİÇEK	143
3. EFFECT OF WHEATGRASS (<i>Triticum aestivum</i> L.) JUICE ON SEEDLING GROWTH AND <i>Rhizoctonia solani</i> ON CORN - İlknur AKGÜN, Rabia AYATA, Ruziye KARAMAN, Gürsel KARACA	149
4. DETERMINATION OF YELLOW RUST DISEASE (<i>Puccinia striiformis</i> f. sp. <i>tritici</i>) RESISTANT OF THE WHEAT LANDRACE COLLECTED FROM ISPARTA AND BURDUR PROVINCES - Demet ALTINDAL, İlknur AKGÜN, Hülya ÖZGÖNEN ÖZKAYA	155
5. USING DIFFERENT METHODS OF ADDING HERBICIDES IN CONTROLLING WATER HYACINTH (<i>Eichhornia crassipes</i>) AND REDUCE IN WATER ENVIRONMENT POLLUTION - Adnan AL-WAGGA, Omar AL-GBURI	162
6. ANTIFUNGAL ACTIVITY OF NANO CALCIUM POLYSULFIDE AGAINST PATHOGENIC FUNGI ON TOMATO - Şerife Evrim ARICI, Ahmet ŞİMSEK, Mustafa ASKIN	170

7. EVALUATION OF CYTOTOXICITY OF THE HERBICIDE GALIGAN 240 EC TO PLANTS - Elena BONCIU	175
8. GROWTH AND GRAIN YIELD PARAMETERS OF SINGLE-PLANTED AND IN-CANOPY GROWN WHEAT (<i>Triticum aestivum</i> L.) - Uğur ÇAKALOĞULLARI, Gülден Deniz ATEŞ ATASOY, Deniz İŞTİPLİLER, Özgür TATAR	179
9. QUALITATIVE MODIFICATIONS PRODUCED IN FEED OF <i>Festuca rubra</i> L. AND <i>Agrostis capillaries</i> L. UNDER INFLUENCE OF UAN LIQUID FERTILIZER - Mirela CIREBEA, Ioan ROTAR, Roxana VIDICAN, Anca PLEŞA, Ovidiu RANTA	184
10. ALLELOPATHY AND ALLELOCHEMICAL INTERACTIONS AMONG PLANTS - Ramona COTRUȚ	188
11. INVESTIGATION ON THE YIELD AND GRAIN QUALITY OF COMMON WHEAT (<i>T. aestivum</i> L.) CULTIVARS GROWN UNDER THE AGROECOLOGICAL CONDITIONS OF CENTRAL BULGARIA - Vania DELIBALTOVA, Manol DALLEV	194
12. APPLICATION OF THE COEFFICIENT OF USING THE ACTIVE SUBSTANCE FOR THE EVALUATION OF THE EFFECT OF CHANGES OF NOZZLE OPERATING PARAMETERS ON THE SPRAYING PROCESS - Katarzyna DEREŃ, Antoni SZEWCZYK, Deta ŁUCZYCKA, Beata CIENIAWSKA	199
13. EVALUATION OF SOME MORPHOLOGICAL, CHEMICAL PARAMETERS AND ANTIOXIDANT CAPACITY OF POMEGRANATE - Cristina DIȚESCU, Narcisa BĂBEANU, Sultana NIȚĂ, Ovidiu POPA	205
14. EFFECT OF BIOCYCLIC HUMUS SOIL ON YIELD AND QUALITY PARAMETERS OF SWEET POTATO (<i>Ipomoea batatas</i> L.) - Lydia Dorothea EISENBACH, Antigolena FOLINA, Charikleia ZISI, Ioannis ROUSSIS, Ioanna TABAXI, Panayiota PAPASTYLIANOU, I. KAKABOUKI, Aspasia EFTHIMIADOU, Dimitrios J. BILALIS	210
15. DETERMINATION OF THRESHING PERFORMANS OF NEW DESING THRESHING UNIT FOR SAGE - Mehmet Emin GOKDUMAN, Deniz YILMAZ	218
16. DETERMINATION OF THE EFFECTS OF GENOTYPE AND SOWING TIME ON PLANT GROWTH AND DEVELOPMENT IN CAMELINA [(<i>Camelina sativa</i> L. (Crantz))] - Merve GÖRE, Orhan KURT	223
17. WEEDS MAPPING FROM WHEAT CROPS - Marga GRADILA, Daniel JALOBA	227
18. ANALYSIS OF SOME QUALITY COMPONENTS TO FEW AMPHIDIPOID LINES OF WHEAT - Paula IANCU, Marin SOARE, Ovidiu PĂNIȚĂ	234
19. DRY MATTER YIELD AND DIGESTIBILITY OF SECOND CROP SILAGE CORN CULTIVATED AFTER CEREALS UNDER ESKISEHIR ECOLOGICAL CONDITIONS - Onur ILERI, Suleyman AVCI, Ali KOC	240
20. RESULTS REGARDING YIELD COMPONENTS AND GRAIN YIELD AT SUNFLOWER UNDER DIFFERENT ROW SPACING AND NITROGEN FERTILISATION CONDITIONS - Viorel ION, Adrian Gheorghe BĂȘA, Marin DUMBRAVĂ, Lenuța Iuliana EPURE	247
21. VARIATION OF CURRENT MORPHOLOGICAL CHARACTERS IN WINTER BARLEY, <i>Hordeum vulgare</i> L. - Nicolae IONESCU, Mihaela Ioana GEORGESCU, Aurel PENESCU, Elena SĂVULESCU, Maria VOICA, Alexandru LAZĂR	255
22. SALINITY EFFECTS ON SWEET CORN YIELD AND WATER USE EFFICIENCY UNDER DIFFERENT HYDROGEL DOSES - Sema KALE, Burcu ARICAN	263

23. EFFECTS OF DIFFERENT SEED SIZES AND SHAPES ON FORAGE YIELD AND QUALITY OF FODDER MAIZE - Emre KARA, Mustafa SÜRME	267
24. EFFECTS OF NANO SULFUR (S) APPLICATIONS ON YIELD AND SOME YIELD PROPERTIES OF BREAD WHEAT - Muharrem KAYA, Ruziye KARAMAN, Aykut ŞENER	274
25. EFFECT OF EXTRA POTASSIUM SUPPLY ON AMINOACID COMPOSITION OF CORN SEED UNDER THE DEFICIT IRRIGATION CONDITIONS (SECTION C) - Yakup Onur KOCA	280
26. PHOSPHINE RESISTANCE OF RUSTY GRAIN BEETLE <i>Cryptolestes ferrugineus</i> (<i>Coleoptera: Laemophloeidae</i>) POPULATIONS IN TURKEY - Erhan KOÇAK, Abdullah YILMAZ, Y. Nazım ALPKENT, Serdar BİLGİNTURAN	286
27. IMPACT OF SOIL TILLAGE REDUCTION ON CULTIVATION PRODUCTIVITY OF WHEAT, BARLEY AND SOYBEAN - Igor KOVAČEV, Nikola BILANDŽIJA, Krešimir ČOPEC, Dubravko FILIPOVIĆ	291
28. A BIBLIOMETRIC REVIEW OF RESEARCH TRENDS IN SALICYLIC ACID USES IN AGRICULTURAL AND BIOLOGICAL SCIENCES: WHERE HAVE BEEN STUDIES DIRECTED? - Muhittin KULAK	296
29. DETERMINATION OF YIELD AND YIELD COMPONENTS OF SOME CRAMBE GENOTYPES IN THE WORLD CRAMBE COLLECTION - Orhan KURT, Tuba ÖZYILMAZ, Merve GÖRE	304
30. IMPROVE OF GRAIN YIELD AND QUALITY OF WINTER WHEAT BY NITROGEN INPUTS - Roxana Maria MADJAR, Gina VASILE SCĂEȚEANU, Andreea ANTON	310
31. CONTROL OF THE CARROT CYST NEMATODE <i>Heterodera carotae</i> BY TANNIN AQUEOUS SOLUTIONS - Lara MAISTRELLO, Nicola SASANELLI, Giacomo VACCARI, Ion TODERAS, Elena IURCU-STRAISTARU	316
32. STUDY OF <i>Rhinoncus pericarpus</i> (Linneus, 1758) (<i>Coleoptera: Curculionidae</i>) BIOLOGY, AN IMPORTANT PEST OF HERB PATIENCE AND RHUBARB IN ROMANIA - Traian MANOLE	322
33. FIRST RECORDS OF NATURAL ENEMIES OF KERMES HERMONENSIS SPODEK & BEN-DOV (<i>Hemiptera: Sternorrhyncha: Kermesidae</i>) IN TURKEY - Hasan MARAL, Halil BOLU	334
34. BIOLOGICAL EFFICACY OF SOME SOIL HERBICIDES AT MAIZE (<i>Zea mays</i> L.) - Anyo MITKOV, Mariyan YANEV, Nesho NESHEV, Tonyo TONEV	340
35. PERFORMANCE OF MAIZE VARIETIES (<i>Zea mays</i> L.) WITH DIFFERENT RATES OF NITROGEN FERTILIZER AND COWDUNG IN MUBI, NORTHERN GUINEA SAVANNA, NIGERIA - Mustapha Alhaji MUHAMMAN, Ruth SAMUEL	346
36. THE EFFECTS OF TILLAGE METHODS AND PLANT DENSITY ON GROWTH, DEVELOPMENT AND YIELD OF SOYBEAN [<i>Glycine max</i> (L.) Merrill] GROWN UNDER MAIN AND SECOND CROPPING SYSTEM: II. GROWTH-DEVELOPMENT COMPONENT – Ferhat ÖZTÜRK, Tahsin SÖGÜT	353
37. THE CHANGE OF SHEAR FORCE AND ENERGY OF COTTON STALK DEPEND ON KNIFE TYPE AND SHEAR ANGLE - F. Göksel PEKİTKAN, Reşat ESGİCİ, A. Konuralp ELİÇİN, Abdullah SESSİZ	360

38. COMPARATIVE STUDY REGARDING THE INFLUENCE OF HERBICIDES ON THE YIELD OF SUNFLOWER CROPS, THE CROPS BEING OBTAINED WITH CONVENTIONAL, CLEARFIELD AND EXPRESSUN TECHNOLOGIES IN THE FIELD CONDITIONS OF MOARA DOMNEASCA - Aurelian PENESCU, Mariana BRAN, Mihaela NICHITA, Nicolae IONESCU, Dumitru Ilie SĂNDOIU, Costică CIONTU, Tudor ȘCHIOPU, Mihai GÎDEA, Cosmin ȘONEA, Ciprian BOLOHAN ...	367
39. DETERMINATION OF THE EFFECTS OF DIFFERENT FERTILIZER APPLICATIONS ON SWEET CORN - Zanko Othman RASHID, Cagatay TANRIVERDI	375
40. EFFECT OF HIGH DILUTIONS OF SODIUM CHLORIDE SOLUTIONS ON WHEAT GERMINATION - PRELIMINARY STUDY - Ieana RÎNDAȘU, Roxana CICEOI, Elena Ștefania IVAN, Florin STĂNICĂ.....	380
41. THE INFLUENCE OF THE INTERACTION OF SOME MINERAL FERTILIZERS ON THE ACCUMULATION OF SOME NUTRITIVE ELEMENTS IN WHEAT GRAINS - Elena ROȘCULETE, Cătălin Aurelian ROȘCULETE	386
42. FACTORS AFFECTING ENERGY CONSUMPTION IN HAMMER MILLS - Hüseyin SAUK, Kemal Çağatay SELVİ	392
43. EFFECT OF LOW TEMPERATURE ON DIAPAUSE EGGS OF <i>Dysdercus cingulatus</i> (Hemiptera: Pyrrhocoridae) - Shahjahan SHAIKH	397
44. RESEARCH AT NIRDPSB BRASOV ABOUT <i>IN VITRO</i> BEHAVIOR OF POTATO PLANTLETS BELONGING TO NEW AMELIORATED LINES AT SRDP TARGU SECUIESC - Andreea TICAN, Carmen Liliana BĂDĂRĂU, Anca BACIU,, Mihaela CIOLOCA	401
45. THE EFFECT OF NITROGEN FERTILIZER ON THE YIELD AND QUALITY IN THE SWEET MAIZE - Mevlüt TÜRK, Mehmet ALAGÖZ	408
46. THE BIOCHEMICAL METHANE POTENTIAL OF <i>Miscanthus giganteus</i> BIOMASS UNDER THE CONDITIONS OF MOLDOVA - Victor ȚÎȚEI	412
47. INDUCING TRANSIENT GENE EXPRESSION IN <i>Nicotiana tabacum</i> PLANT BY AGROINFILTRATION METHOD - Anca Amalia UDRIȘTE	416
48. THE ROLE OF BUFFER ZONES IN ENSURING THE COEXISTENCE OF GM AND NON-GM MAIZE - Viorica URECHEAN, Dorina BONEA	420

MISCELLANEOUS

1. COMPARISON BETWEEN PRISTINE PURE-BEECH STAND AND MIXED BEECH-OAK STAND USING AN UNMANNED AERIAL VEHICLE - Tiberiu Paul BANU, Manfred SCHÖLCH, Constantin BANU, Gheorghe Florian BORLEA.....	429
2. MODAL ANALYSIS OF FIELD SPRAYER BOOM DESIGN FOR DIFFERENT MATERIALS - Ali BAYAT, Medet İTMEÇ, Ali BOLAT	434
3. EVALUATION OF TILLAGE SYSEMS ON SOIL FUNGUS MICROFLORA UNDER WINTER WHEAT CULTIVATION - Serkan BAYMAN, M. Murat TURGUT	438
4. TOTAL CONTENT OF SOME ANTIOXIDANTS IN TEN ROMANIAN POTATO GENOTYPES - Carmen Liliana BĂDĂRĂU, Maria Floriana ȘTEFAN, Cristina Maria CANJA, Mirabela Ioana LUPU	442

5. ESTIMATION OF THE LIQUID COVER OF SELECTED DICOTYLEDONOUS PLANTS IN VARIOUS PHASES OF DEVELOPMENT DURING SPRAYING - Beata CIENIAWSKA, Deta LUCZYCKA, Antoni SZEWCZYK, Katarzyna DEREŃ	448
6. DIVERSITY, DISTRIBUTION AND ECOLOGY OF THE FOREST NATURAL HABITATS IN THE BRATOVOEȘTI FOREST, DOLJ COUNTY - Florin Dorian COJOACĂ, Mariana NICULESCU	453
7. PHYTOCHEMICAL RESEARCH ON AERIAL PARTS OF <i>Raphanus raphanistrum</i> subsp. <i>landra</i> (Moretti ex DC.) Bonnier & Layens - Aurora DOBRIN, Vlad Ioan POPA, Constantin Daniel POTOR, Mihaela Ioana GEORGESCU	458
8. WHICH SHRUB SPECIES SHOULD BE USED FOR THE ESTABLISHMENT OF FIELD SHELTERBELTS IN ROMANIA? - Cristian Mihai ENESCU	464
9. MORPHOLOGICAL ASPECTS ABOUT GERMINATION OF <i>Vulpia myuros</i> (L.) C.C. Gmel. CARYOPSIS - Mihaela Ioana GEORGESCU, Vlad Ioan POPA, Daniel Constantin POTOR, Nicolae IONESCU, Elena SĂVULESCU	470
10. MELLIFEROUS POTENTIAL OF SILVER LINDEN TREES (<i>Tilia tomentosa</i> Moench.) GROWING IN THE FORESTS FROM SOUTH ROMANIA - Nicoleta ION, Răzvan COMAN, Viorel ION	474
11. INVESTIGATION OF DIFFERENT TILLAGE AND SEEDING METHODS IN SAFFLOWER (<i>Carthamus tinctorius</i> L.) CULTIVATION - Hüseyin KÜÇÜK, Davut AKBOLAT	481
12. NEW INSIGHTS INTO THE MULTIPLE PROTECTIVE FUNCTIONS OF DIATOMACEOUS EARTH DURING STORAGE OF AGRICULTURAL PRODUCTS - Carmen LUPU, Elena DELIAN, Lenuța CHIRA, Adrian CHIRA	487
13. BIOMASS QUALITY OF SOME <i>Poaceae</i> SPECIES AND POSSIBLE USE FOR RENEWABLE ENERGY PRODUCTION IN MOLDOVA - Ivan MUNTEAN, Victor ȚÎȚEI, Andrei GUDIMA, Andrei ARMAȘ, Mihai GADIBADI	497
14. RESEARCH OF THE COLEOPTERA (<i>Cerambycidae</i> and <i>Lucanidae</i>) FOUND IN THE NATURAL HABITATS OF THE GATEJESTI-BUNESTI FOREST - Laurențiu NICULESCU, Ion MITREA	503
15. STUDIES ON THE VITALITY AND THE STATE OF HEALTH OF CHARACTERISTIC SPECIES OF TREES FROM THE FOREST HABITATS FOUND IN THE PROTECTED AREA ROSCI 0128 (NORDUL GORJULUI DE EST) - Mariana NICULESCU, Liviu Aurel OLARU, Silvestru Ilie NUȚĂ	507
16. DEVELOPMENT OF TAILOR-MADE FOOD WASTE PREVENTION MEASURES BASED ON CONSUMER TYPE ANALYSIS - Sandra SCHWÖDT, Gudrun OBERSTEINER	513
17. PRECISION OF DROUGHT BASED ON THE TOPSIS METHOD - Çağatay TANRIVERDİ, Hasan DEĞİRMENCİ, Mahmut TEKİNERDOĞAN, Engin GÖNEN, Firat ARSLAN, Atılğan ATILGAN	516

SOIL SCIENCES

THE EFFECTS OF DIFFERENT ZN DOSES AND MYCORRHIZAE APPLICATION ON HORSE BEAN GROWTH AND NUTRIENT UPTAKE UNDER STERILE AND NON STERILE SOIL CONDITIONS

Cagdas AKPINAR¹, Ibrahim ORTAS², Ahmet DEMIRBAS³

¹Department of Organic Farming Business Management, Kadirli School of Applied Sciences, Osmaniye Korkut Ata University, Osmaniye, Turkey

²Department of Soil Science and Plant Nutrition, Faculty of Agriculture, Cukurova University, Adana, Turkey

³Department of Crop and Animal Production, Vocational School of Sivas, Cumhuriyet University, Sivas, Turkey

Corresponding author email: cagdasakpinar@osmaniye.edu.tr

Abstract

Due to semi-arid Mediterranean climate conditions and soil properties, south east Turkey's soils usually do not contain sufficient amount of plant nutritious elements, especially phosphorous (P) and zinc (Zn). This scarcity conditions is one of the factors limiting agriculture productivity and fertility and consequently, decrease the crop production. Mycorrhiza can be able to increase soil fertility and enhance nutrient uptake and plant growth. The aim of this study was to investigate the effects of mycorrhizal inoculation and different Zn doses on horse bean (*Vicia faba* L.) growth and nutrient uptake under sterile and non-sterile soils conditions. The experiment was carried out under greenhouse conditions with three replications. Horse bean was used as a test plant and *Funneliformis mosseae* was used as mycorrhiza species (1000 spore plant⁻¹). Sterile and non-sterile Karaburun soils series from Çukurova region were used. Two doses of Zn (0 and 5 mg kg⁻¹) were used as fertilizer.

In both sterilized, non-sterilized soils mycorrhizal inoculation increased shoot dry matter and nutrient concentration. The results shown that shoot dry matter and nutrient concentration significantly increased of horse bean plants in non-sterilized soils. Maximum value of % P and Zn concentration was determined and in non-sterilized soils mycorrhiza inoculated have 0.28 % P (0 mg kg⁻¹Zn) and in mycorrhiza inoculated sterile soils, 52.1 mg kg⁻¹Zn (5 mg kg⁻¹Zn). The results are encouraging that the mycorrhiza can be used as an organic fertilizer for plant production under large arable conditions.

Key words: mycorrhizae, horse bean, zinc, phosphorus.

INTRODUCTION

Middle East-originated horse bean (*Vicia faba* L.) is among the earliest cultured legumes after chickpea and pea. Horse bean is commonly grown for human nutrition, but it is also used as animal feed (Prolea, 2014). Turkey, especially the Mediterranean Region, has loamy soils with high clay and low organic matter contents because of climate zone and geographical location. Undesired physical, chemical and biological soil characteristics also reduce available plant nutrient concentrations. Soils of Çukurova Region (Eastern Mediterranean Region) with intensive agricultural practices experience nutrient deficiency place to place mostly based on soil parent material and especially phosphorus (P) and zinc (Zn) deficiencies are the common cases in these soils (Ibriki, 1994).

Mycorrhizal symbiosis between plants and fungi has been one of the most well-known plant-fungus associations since about 450 million years and is of significant importance for growth and persistence of terrestrial plants (Redecker et al., 2000). Obligate symbiotic arbuscular mycorrhizal fungi (AMF) belongs to endomycorrhiza group of *Glomeromycota* family and the fungi is in symbiotic association with about 80% of the plant species (Smith and Read, 2008). Horse bean is also among these plant species which is significantly mycorrhizal dependent (Ortas, 2012a). Turkey has also grown horse bean which can form associations with mycorrhizal fungi (Kaschuk et al., 2010). Plant roots are not the only dominant factors in plant nutrient uptake, but there are some other factors effective in nutrient uptake processes of the plants. It was reported in previous studies that hyphae-producing fungi species, also called

as mycorrhiza, which can be seen under microscope and have symbiotic associations with plant roots, may also provide significant supports in plant nutrient uptake from the soils (Ryan et al., 2008). Arbuscular mycorrhiza fungi (AMF) aid in enlargement of root surface area in rhizosphere area and alter rhizosphere pH, thus increase P uptake of plants from the soils (Ortas and Rowell, 2004; Ortas and Akpinar, 2011). In case of an efficient infection between the mycorrhiza and the plants, plants give 5-10% of the carbon to mycorrhiza fungi and in return mycorrhiza facilitates plant nutrient flow (Hodge and Storer, 2015) and water uptake (Augé, 2001). It has been found that mycorrhizal inoculation increased plant especially P, Zn and Cu concentrations when effective root infection occurs (Smith and Read, 2008). Besides plant growth, AMF has several positive impacts on soil quality (Celik et al., 2011).

Zn uptake mechanism of mycorrhiza is similar with the phosphorus uptake mechanism and supplies about 60% of the Zn provided to plants through mycorrhizal hyphae from outside the root region (rhizosphere) (Ortas et al., 2011). Arbuscular mycorrhiza not only facilitates P uptake of the host plant, but also significantly increase Zn uptake of the plants (Ortas and Akpinar, 2011; Ortas, 2012b). Therefore, existence of mycorrhiza fungi hyphae is a quite significant issue for the uptake of slowly available plant nutrients like Zn. According to Thompson (1990), non-existence of Zn deficiency of the plant species grown in soils with low DTPA-extractable Zn concentrations mostly resulted from Zn supplied from the soils through AM.

Zn is a quite significant microelement for human health. Zn deficiency is experienced about 60% of Turkish soils (Eyüpoğlu et al., 1996), therefore mycorrhiza utilization in agricultural soils is a significant issue to prevent Zn deficiency. Mycorrhiza utilization may have a strategic importance especially in soils of inner semi-arid regions. This study was conducted to investigate the effects of different Zn doses and mycorrhiza inoculations on

nutrient uptake and plant growth of horse beans grown in sterilized and unsterilized soils poor in plant nutrients.

MATERIALS AND METHODS

The experiment was conducted on Karaburun soil series (*Typic Xerorthent*) which is nutrient deficient (Table 1). Soil used in the investigation which is silt clay loam, slightly alkaline (pH 7.4), calcareous (23.2%), phosphorus content low (26.0 kg·ha⁻¹), and potassium content is sufficient (630 kg·ha⁻¹). The half of the soil was sterilized by using an autoclave at 120°C for two hours. Two levels of Zn, 0 and 5 mg·kg⁻¹ soil, as zinc sulfate (ZnSO₄·7H₂O) were applied.

Plants were inoculated with the AMF *Funneliformis mosseae*. A level of 1000-spore per pot was placed 3 cm below the seeds. The non-inoculated pots received the same amount of mycorrhizal spore-free inoculum.

Horse bean (*Vicia faba* L.) was used as a test plant. In the study, plastic pots with a capacity of 3 kg were used and 5 seeds were planted at the beginning of each pot. After germination, they were reduced to 3 plants. Distilled water was added daily to maintain the moisture at 75% of field capacity.

At the harvest of each pot, total plant biomass (dry weight of root and shoot) were recorded. Plant materials, shoots and roots were washed thoroughly with distilled. Then plants were oven-dried at 65°C for 48 h. The dry material was ground using a Tema mill, and 0.2 g of the ground plant material was ashed at 550°C, then dissolved in 3.3% HCl. The concentration of phosphorus was determined according to Murphy and Riley (1962). An ICP-spectrophotometer (ICP-OES; Thermo ICAP7000) was employed to determine the concentration of K, Zn and Fe in the plant samples. The root staining procedure described by Koske and Gemma (1989). The percentage of AMF colonization was identified by method of Giovanetti and Mosse (1980).

Table 1. Some physical and chemical properties of Karaburun series soils used in experiments

Soil	P ₂ O ₅	K ₂ O	Fe	Zn	Cu	Mn	pH	CaCO ₃	Texture
Depth	kg·ha ⁻¹		mg·kg ⁻¹			%			
(0-30)	26.0	630.0	3.02	0.38	0.57	3.21	7.41	23.2	SiCL

Mycorrhizal Dependency (MD)

After harvest, the total dry weight of the seedlings was recorded and inoculation effectiveness of the seedlings by AMF was calculated using the following formula by Ortas (2012a):

$$(MD) = \frac{TDW (+M) - TDW (-M)}{TDW (+M)} \times 100$$

TDW= total dry weight; +M, inoculated plant; -M, non-inoculated plant

Statistical Analysis

All statistical analyses (Tukey test and correlation) were performed using the SPSS 22.0 for Windows computer program.

RESULTS AND DISCUSSIONS

Effects of different Zn doses and mycorrhiza inoculation on shoot (SDW) and root dry weight (RDW) production of horse bean were investigated and results are presented in Table 2. Present findings revealed that mycorrhiza inoculations significantly increased both shoot and root dry matter production. While shoot dry matter production was 7.62 g pot⁻¹ and 7.44 g pot⁻¹ in unsterilized and sterilized control treatment respectively (Table 2). With 5 mg kg⁻¹Zn and mycorrhiza treatments, SDW was 8.36 g pot⁻¹ under sterilized and 8.08 g pot⁻¹

under the sterilized soil conditions. The greatest shoot dry matter production was observed as 9.18 g pot⁻¹ in mycorrhiza-inoculated and non-Zn-treated sterilized control soil. Similar findings were also observed in root dry matter productions. Mycorrhiza had positive impacts on plant growth and development under both sterilized and unsterilized soil conditions. Considering the general averages, it was observed that the greatest shoot dry matter production of horse bean plants in unsterilized soils was 8.21 g pot⁻¹, the greatest root dry matter production was observed as 7.63 g pot⁻¹ in sterilized soils. Similarly, Akpınar (2011) also reported that AM inoculations significantly increased dry matter production of horse bean plants. Also Ortas (2012a) showed that during 1997 to 1999 mycorrhizal inoculation significantly increased the horse bean yield under field conditions. Following the soil sterilization, since the other microorganisms, existing in soil and competing with plant roots for nutrients, were removed from the soil, mycorrhiza fungi had much more impacts on plant growth and development. Indigenous mycorrhiza also exist in unsterilized soils to some extent and thus had positive impacts on plant growth and development also in unsterilized soils (Ortas, 2003).

Table 2. Effects of different zinc doses and mycorrhiza inoculation on shoot and root dry matter production of horse bean plants grown in sterilized and unsterilized soils

		Shoot Dry Weight		Root Dry Weight	
		g pot ⁻¹			
(-) Sterile	(-M)	(0 mg kg ⁻¹ Zn)	7.62 ±0.5 ^b	6.94 ±0.0 ^{ab}	
		(5 mg kg ⁻¹ Zn)	7.68 ±0.1 ^{ab}	5.71 ±0.6 ^b	
		(0 mg kg ⁻¹ Zn)	9.18 ±0.9 ^a	7.61 ±0.2 ^{ab}	
		(5 mg kg ⁻¹ Zn)	8.36 ±0.0 ^{ab}	9.56 ±1.3 ^{ab}	
		General Average	8.21^A	7.45^A	
(+) Sterile	(-M)	(0 mg kg ⁻¹ Zn)	7.44 ±0.2 ^b	7.36 ±0.1 ^{ab}	
		(5 mg kg ⁻¹ Zn)	7.81 ±0.4 ^{ab}	6.27 ±0.1 ^{ab}	
		(0 mg kg ⁻¹ Zn)	8.69 ±0.0 ^{ab}	9.88 ±2.5 ^{ab}	
		(5 mg kg ⁻¹ Zn)	8.08 ±0.1 ^{ab}	7.02 ±0.1 ^a	
		General Average	8.00^B	7.63^A	

± Standard error, P<0.05

The greatest root infection was observed as 65% in 5 mg kg⁻¹ Zn-treated and mycorrhiza-inoculated sterilized soils (Table 3). The reason of that high infection rate was considered as absence of other organisms to compete with mycorrhiza. The mycorrhizal inoculum increased the root colonization of horse bean plants compared with the non-inoculated treatments (Ortas, 2012a). Similar with the present findings, previous researchers also reported that Zn fertilization did not influence mycorrhiza infections (Ortas et al., 2002; Subramanian et al., 2009).

Mycorrhizae dependency (MD) was calculated after harvest. As can be seen in the table, MD increased with non-sterile soil. At non sterile soil was 19.31 %, and sterile soil was 13.50 % as calculated (Table 3). This means that mycorrhiza inoculation at the sterile soil was

not sufficient. Also, mycorrhizal dependency was found higher in Zn applications as 25.29 %. Ortas (2012a) found that horse bean showed the highest mycorrhizal dependency in all 3 years, especially in P0 plots under field conditions. Tawaraya (2003) reported that mean MD values of 44% for field crops (37 species).

When the present findings were assessed for root infection and mycorrhizal dependence, root infections were observed in control treatments of sterilized soils even at quite low levels because of the existence of indigenous mycorrhiza. Since non-sterile soils have indigenous mycorrhizae which may significantly increase plant growth. As can be seen in Table 2, there is small differences in dry weight in between sterile and non-sterile soil treatments.

Table 3. Effects of different zinc doses and mycorrhiza inoculation on root infection and mycorrhizal dependence of horse bean plants grown in sterilized and unsterilized soils

			Root Infection %	Mycorrhizal Dependency %
(-) Sterile	(-M)	(0 mg kg ⁻¹ Zn)	10 ±0.0 ^c	
		(5 mg kg ⁻¹ Zn)	5 ±7.1 ^c	
	(M)	(0 mg kg ⁻¹ Zn)	15 ±7.1 ^c	13.33
		(5 mg kg ⁻¹ Zn)	15 ±7.1 ^c	25.29
	General Average		11.3 ^B	19.31
(+) Sterile	(-M)	(0 mg kg ⁻¹ Zn)	0 ±0.0 ^c	
		(5 mg kg ⁻¹ Zn)	0 ±0.0 ^c	
	(M)	(0 mg kg ⁻¹ Zn)	40 ±0.0 ^b	20.30
		(5 mg kg ⁻¹ Zn)	65 ±7.1 ^a	6.71
	General Average		26.3 ^A	13.50

± Standard error, P<0.05

When the present findings were assessed for plant nutrients, it was observed that while the lowest P concentration was 0.13% in 5 mg kg⁻¹ Zn-treated and mycorrhiza-inoculated sterile soils and the greatest P concentration was 0.28% in mycorrhiza-inoculated unsterilized soils (Table 4). Although the differences between sterilized and unsterilized soils were not found to be significant, P concentrations generally increased with mycorrhiza inoculations as compared to control treatments. It was reported that mycorrhiza fungi significantly increased the uptake of slowly available nutrients, especially the phosphorus under controlled conditions (Grant et al., 2005; Ortas and Akpinar, 2006; Ortas, 2012b). Similar with P concentrations, plant tissue potassium (K) concentrations also

increased with mycorrhiza inoculations. With regard to Fe concentrations, the greatest value was observed as 308.4 mg kg⁻¹ in 5 mg kg⁻¹Zn-treated and mycorrhiza-inoculated unsterilized soils. The lowest Zn concentration (26.7 mg kg⁻¹Zn) was observed in sterilized and non-mycorrhiza-inoculated control treatments and the greatest Zn concentration (52.1 mg kg⁻¹Zn) was observed in 5 mg kg⁻¹Zn-treated and mycorrhiza-inoculated sterilized soils. Previous researchers also indicated that mycorrhiza treatments increased uptake of immobile nutrients, especially Zn, at acceptable levels (Ortaş and Akpinar, 2006; Smith and Read, 2008). In another study, *Fu. mosseae* inoculations increased nutrient uptake levels of horse bean plants (El-Sayad et al., 2002). Ortas (2012a) found that mycorrhizal inoculation

significantly increased the horse bean P and Zn concentrations in three years under field conditions. With regard to general averages,

the plants grown in unsterilized soils had higher nutrient concentrations than the plants grown in sterilized soils.

Table 4. Effects of different zinc doses and mycorrhiza inoculation on P, K, Fe and Zn concentration of horse bean plants grown in sterilized and unsterilized soils

		P	K	Fe	Zn	
		%		mg kg ⁻¹		
(-) Sterile	(-M)	(0 mg kg ⁻¹ Zn)	0.17 ±0.1 ^a	2.5 ±0.4 ^a	147.3 ±16.5 ^{bc}	36.0 ±0.0 ^{cd}
		(5 mg kg ⁻¹ Zn)	0.22 ±0.0 ^a	3.1 ±0.0 ^a	292.8 ±4.2 ^a	35.7 ±2.2 ^{cd}
	(+M)	(0 mg kg ⁻¹ Zn)	0.28 ±0.1 ^a	3.3 ±0.4 ^a	307.8 ±12.7 ^a	46.7 ±3.3 ^{ab}
		(5 mg kg ⁻¹ Zn)	0.23 ±0.1 ^a	3.9 ±0.9 ^a	308.4 ±62.6 ^a	41.0 ±3.0 ^{ac}
General Average			0.23 ^A	3.20 ^A	264.1 ^A	39.9 ^A
(+) Sterile	(-M)	(0 mg kg ⁻¹ Zn)	0.15 ±0.0 ^a	2.6 ±0.2 ^a	153.7 ±57.7 ^{bc}	26.7 ±1.2 ^d
		(5 mg kg ⁻¹ Zn)	0.13 ±0.0 ^a	3.3 ±0.3 ^a	255.3 ±6.7 ^{ab}	33.5 ±1.5 ^{cd}
	(+M)	(0 mg kg ⁻¹ Zn)	0.19 ±0.0 ^a	3.1 ±0.3 ^a	254.2 ±18.6 ^{ab}	41.9 ±3.6 ^{ac}
		(5 mg kg ⁻¹ Zn)	0.25 ±0.0 ^a	3.7 ±0.1 ^a	118.9 ±1.6 ^c	52.1 ±4.7 ^a
General Average			0.18 ^B	3.18 ^A	195.6 ^B	38.6 ^A

± Standard error, P<0.05

The correlation analysis among the traits evaluated in this study is given in Table 5. As seen from the table, a positive relationship was found between shoot dry weight, P and Zn also between root dry weight and mycorrhizal dependency. Previous studies have found similar correlations between certain plant

developmental parameters and mycorrhizal dependency (Azcon and Ocampo, 1981; Chalk et al., 2006). In the present study, Zn concentration was positively correlated with shoot dry weight, P, K concentration, root infection and negatively correlated with mycorrhizal dependency (Table 5).

Table 5. Correlation values of parameters evaluated in the study

Parameters	SDW	RDW	P	K	Fe	Zn	Inf.	MD
SDW	1	.430	.516*	.293	.474	.641**	.350	.098
RDW		1	.284	.409	.272	.277	.270	.717*
P			1	.542*	.333	.657**	.379	-.296
K				1	.431	.584*	.340	.074
Fe					1	.076	-.357	.716*
Zn						1	.764**	-.842**
Infection							1	-.650
MD								1

*: P <0.1, **: P < 0.05

SDW: Shoot Dry Weight; RDW: Root Dry Weight; P: Phosphorus; K: Potassium; Fe: Iron, Zn: Zinc; Inf: Root Infection; MD: Mycorrhizal Dependency

CONCLUSION

Present findings revealed that AMF, a beneficial fungi species and already existing in soils, had positive impacts on growth, development and nutrient uptake of horse bean plants. Such findings clearly indicated that mycorrhiza inoculations could play significant roles in growth and development of horse bean and similar plants grown especially in soils poor in plant nutrients. Also the results are shown that indigenous mycorrhizae have

significant effect on plant growth and nutrient uptake as well. In this sense, agricultural strategies should be developed for naturally existing mycorrhiza in soils and thus for sustainable agricultural practices. Use of such natural rhizosphere mechanisms in agricultural practices will provide significant contributions to both the ecosystem and the economy of the countries largely dependent on foreign countries for chemical fertilizers. For sustainable agriculture, use of mycorrhiza as a biotical fertilizer will also provide significant

contributions in reducing chemical fertilizer-induced environmental problems. Use of mycorrhiza in agricultural practices could also be recommended for healthy nutrition, food safety and quality.

REFERENCES

- Akpınar Ç., 2011. Kanola sonrası yetiştirilen II. ürün mısır bitkisine mikoriza aşılamaşının verim ve besin elementleri alınma etkisi. Çukurova Üniversitesi, Fen Bil. Enstitüsü. Doktora Tezi. Adana.
- Augé R.M., 2001. Water relations, drought and vesicular-arbuscular mycorrhizal symbiosis. *Mycorrhiza*, 11:3-42.
- Azcon R., Ocampo J.A., 1981. Factors affecting the vesicular arbuscular infection and mycorrhizal dependency of thirteen wheat cultivars. *New Phytol*, 87: 677-685.
- Celik I., Barut Z.B., Ortas I., Gok M., Demırbas A., Tulun Y., Akpınar C., 2011. Impacts of different tillage practices on some soil microbiological properties and crop yield under semi-arid Mediterranean conditions. *International Journal of Plant Production*, 5 (3): 237-254.
- Chalk P.M., Souza R.F., Urquiaga S., Alves B.J.R., Boddey R.M., 2006. The role of arbuscular mycorrhiza in legume symbiotic performance. *Soil Biol. Biochem*, 38: 2944-2951.
- El-Sayed El-S.A., El-Didamony G., El-Sayed E.F., 2002. Effects of mycorrhizae and chitin-hydrolysing microbes on vicia faba. *World Journal of Microbiology & Biotechnology*, 18:505-515.
- Eyüpoğlu F., Kurucu N., Talaz S., 1996. Türkiye topraklarının bitkiye yararlı bazı mikro elementler (Fe, Cu, Mn, Zn) bakımından genel durumu. *Toprak ve Su Kaynakları Araştırma Yıllığı-1995*, Yayın No: 98, 338-350, Ankara.
- Gioannetti M., Mosse B., 1980. An evaluation of techniques for measuring vesicular-arbuscular mycorrhiza in roots. *New Phytologist*, 84:489-500.
- Grant C., Bittman S., Montreal M., Plenchette C., Morel C., 2005. Soil and fertilizer phosphorus: effects on plant P supply and mycorrhizal development. *Canadian Journal of Plant Sci.*, 85:3-14.
- Hodge A., Storer K. 2015. Arbuscular mycorrhiza and nitrogen: implications for individual plants through to ecosystems. *Plant and Soil*, 386:1-19.
- Ibrikci H., 1994. Macro element status of mandarin orchards in southern Turkey. *Communications in Soil Science and Plant Analysis*, 25(17-18):2791-2980.
- Kaschuk G., Leffelaar P., Giller K., Alberton O., Hungria M., Kuyper T., 2010. Responses of legumes to rhizobia and arbuscular mycorrhizal fungi: a meta-analysis of potential photosynthesis that elimination of symbioses. *Soil Biol Biochem*, 42:125-127.
- Koske R.E., Gamma J.N., 1989. A modified procedure for staining roots to detect VAM. *Mycological Research*, 92:486-505.
- Murphy J., Riley J.P., 1962. A modified single solution method for determination of phosphate in natural waters. *Anal. Chim. Acta*, 27:31-36.
- Ortas I., 2012b. Do maize and pepper plants depend on mycorrhizae in terms of phosphorus and zinc uptake? *J Plant Nutr*, 35 (11):1639-1656.
- Ortas I., 2003. Effect of selected mycorrhizal inoculation on phosphorus sustainability in sterile and non-sterile soils in the Harran Plain in South Anatolia. *J Plant Nutr*, 26 (1):1-17.
- Ortas I., 2012a. The effect of mycorrhizal fungal inoculation on plant yield, nutrient uptake and inoculation effectiveness under long-term field conditions. *Field Crops Research*, 125:35-48.
- Ortas I., Akpınar C., 2006. Response of kidney bean to arbuscular mycorrhizal inoculation and mycorrhizal dependency in P and Zn deficient soils. *Acta Agr Scand B-S P*, 56:101-109.
- Ortas I., Akpınar C., 2011. Response of maize genotypes to several mycorrhizal inoculums in terms of plant growth, nutrient uptake and spore production. *J Plant Nutr*, 34 (7):970-987.
- Ortas I., Ortakci D., Kaya Z., Cinar A., Onelge N., 2002. Mycorrhizal dependency of sour orange in relation to phosphorus and zinc nutrition. *J Plant Nutr*, 25:1263-1279.
- Ortas I., Rowell D.L., 2004. Effect of ammonium and nitrate on indigenous mycorrhizal infection, rhizosphere pH change and phosphorus uptake by sorghum. *Communications in Soil science and plant Analysis*, 35:1923-1944.
- Ortas I., Sarı N., Akpınar C., Yetisir H., 2011. Screening mycorrhiza species for plant growth, P and Zn uptake in pepper seedling grown under greenhouse conditions. *Scientia Horticulturae*, 128: 92-98.
- Ortas I., Akpınar C., 2011. Response of maize genotypes to several mycorrhizal inoculums in terms of plant growth, nutrient uptake and spore production. *Journal of Plant Nutrition*, 34 (7):970-987.
- Prolea, 2014. Lesgraines : pois, féveroles, lupin. Prolea. L'alimentation animal.
- Redecker D., Kodner R., Graham L.E., 2000. Glomalean fungi from the ordovician. *Science*, 289:1920-1921.
- Ryan M.H., Mcinerney J.K., Record I.R., Angus J.F., 2008. Zinc bioavailability in wheat grain in relation to phosphorus fertilizer, crops sequence and mycorrhizal fungi. *Journal of the Science of Food and Agriculture*, 88:1208-1216.
- Smith S., Read D.J., 2008. *Mycorrhizal Symbiosis*. Academic Press, San Diego, Ca.
- Subramanian K.S., Tenshia V., Jayalakshmi K., Ramachandran V., 2009. Role of arbuscular mycorrhizal fungus (*Glomus intraradices*)-(fungusaided) in zinc nutrition of maize. *J Agric Biotechnol Sustain Dev*, 1:029-038
- Tawaraya K., 2003. Arbuscular mycorrhizal dependency of different plant species and cultivars. *Soil Sci. Plant Nutr*. 49(5):655-668.
- Thompson J.P., 1990. Soil sterilization methods to show VA- mycorrhizae aid P and Zn nutrition of wheat in vertisols. *Soil Biology & Biochemistry*, 22:229-240.